



# MILK

## Nutritious by nature

The science behind  
the health and nutritional impact  
of milk and dairy foods

# Cardiovascular disease

The focus on dairy foods and cardiovascular disease (CVD) is often in relation to saturated fat. There is an assumption that because some dairy foods contain saturated fatty acids and dairy in general contributes to saturated fat intake in the diet, that it also increases the risk of cardiovascular disease. Yet the majority of epidemiological studies have shown no adverse effects of regularly consuming milk and dairy foods such as yogurt and cheese on cardiovascular health, irrespective of fat content. In fact, in some cases a cardio-protective effect has been observed.

The explanation for this may lie in the complex composition of milk and dairy foods which, in addition to saturated fat, contain other nutrients and bioactive components such as calcium, potassium and bioactive peptides in the dairy matrix which may be beneficial to cardiovascular health. In addition, the overall fatty acid profile of milk and dairy may not have the detrimental effect on blood lipids or other cardiovascular parameters that has been assumed.

## Observational studies

**Ideally, the relationship between dairy foods and cardiovascular disease would be measured by very large, very long-term intervention trials. In practice, however, the best available data are often from large, long-term observational studies.** There are several

studies of this type in European populations dating back over a number of years. These give an indication of the potential health effects of milk and dairy foods in relation to the particular amounts, types and patterns of consumption in Europe. For example, a prospective cohort study from the UK reports that men who drank the most milk (around a pint / 568ml of whole milk a day) had fewer heart attacks and fewer strokes than those who had little or no milk in their diets<sup>1</sup>. The large Netherlands Cohort Study, consisting of over 120,000 men and women, showed no association between total milk product consumption and stroke mortality, although butter and dairy fat was associated with a small (7%) increased risk of all-cause and heart disease mortality among women<sup>2</sup>. Data from the smaller Dutch Hoorn Study also found that overall dairy intake was not associated with CVD mortality but the intake of high-fat dairy products was<sup>3</sup>. In contrast, the Rotterdam Study reports that in an

older Dutch population, high-fat dairy was associated with a reduced risk of fatal stroke; total dairy consumption or the intake of specific dairy products was not related to CVD events<sup>4</sup>.

In another Dutch prospective cohort, there was again no evidence that dairy products increased risk of heart disease or stroke<sup>5</sup>. In fact, high intakes of total and low-fat dairy were associated with a lower coronary heart disease (CHD) risk in participants without hypertension. Low-fat dairy consumption was also associated with reduced risk of stroke in cohorts of Swedish men and women<sup>6</sup>. In the same cohorts, a high intake of fermented milk (yogurt and cultured sour milk) was found to reduce CVD risk<sup>7</sup>. Fermented milk and cheese were also associated with reduced cardiovascular disease mortality<sup>8</sup>. There was, however, an increased CVD mortality risk reported in this study in those drinking three or more glasses of (non-fermented) milk a day compared with less than one glass. The reason for this discrepancy in the same cohort is not clear and the authors urge a cautious interpretation of the results. Moreover, when these data were re-examined, milk consumption was associated with a lower risk of CVD

mortality<sup>9</sup>. In line with the majority of epidemiological studies, a Danish investigation reports no adverse effects of dairy on cardiovascular health<sup>10</sup>. The French MONICA project conducted over 14 years, found that dairy consumption (particularly milk intake) as part of a diverse, healthy diet was associated with the lowest mortality rate mostly due to reduced cardiovascular deaths<sup>11</sup>. Similarly, in the large French prospective cohort NutriNet-Santé, dairy product consumption was not associated with total CVD or CHD risks, and furthermore consumption of fermented dairy, such as cheese and yogurt, was associated with a 19% reduction in the risk of cerebrovascular disease<sup>12</sup>. The findings of the large cohort PURE-study which included 136,384 individuals from 21 countries and 5 continents, demonstrated that dairy consumption (predominantly milk and yogurt) was associated with lower risk of mortality and major CVD events<sup>13</sup>.

## Meta-analyses for milk

**A number of analyses have pooled the data from individual prospective studies such as these and their results strengthen the evidence that regularly consuming milk and other dairy products does not increase risk of cardiovascular disease and may even have a protective effect**<sup>1,14-26</sup>. In relation to milk, an

overview conducted in 2010 concluded that milk drinking is not harmful and may be associated with a small but worthwhile reduction in risk of coronary heart disease (8%) and a more substantial reduction in stroke risk (21%) for those who drank the most milk compared with those who drank the least<sup>15</sup>. The pooled results of seventeen studies in 2011 also found milk intake was associated with a small potential reduction in overall cardiovascular risk of 6% for each 200ml of milk consumed a day<sup>16</sup>. This analysis found no association between high intakes of either regular-fat or low-fat dairy products and increased risk of death from cardiovascular disease. Similarly, systematic reviews in 2015 and 2017 examining milk consumption and cardiovascular disease mortality observed no consistent association<sup>17,18</sup>. This was also the conclusion for milk and CVD risk in a meta-analysis published in 2016; milk intake was found to be neutral with respect to risk of stroke and coronary artery disease<sup>19</sup>. In a 2021 analysis of six studies comparing the those with the highest regular-fat milk

consumption with the lowest, a greater risk of CHD was observed. However, for milk intake overall, those in the highest consumption group had a reduced risk of ischemic stroke compared with those in the lowest<sup>20</sup>.

## Meta-analyses for dairy products

**Meta-analyses also support neutral or beneficial effects of other dairy foods on cardiovascular disease.**

Twenty-two prospective cohort studies were included in an analysis published in 2015 which examined stroke and CHD incidence in relation to intake of individual dairy foods, and to low- and regular-fat dairy<sup>21</sup>. Cheese consumption was associated with a 16% decreased heart disease risk, and both cheese and low-fat dairy foods were associated with reduced risk of stroke (9% and 7% respectively). An earlier meta-analysis in 2014 looking specifically at stroke also reported similar reductions in risk with low-fat dairy (9%) and cheese intake (6%) and, in addition, with total dairy (12%) and fermented milk (20%)<sup>22</sup>. Similarly, in another meta-analysis including 18 studies which had examined dairy intake and stroke risk, milk and cheese consumption were associated with reduced risk of stroke; risk reductions were maximal around 125g/day for milk (16%) and from 25g/day upwards for cheese (9%)<sup>23</sup>. Cheese was also associated with a lower risk of stroke (13%) in a meta-analysis published in 2016, as was total dairy intake (9%)<sup>24</sup>. In addition, cheese intake was associated with an 18% reduced risk of coronary heart disease. A beneficial effect of cheese was supported by a meta-analysis of prospective cohort studies published in 2017 in which cheese intake was associated with 10%, 14% and 10% reduced risks of total CVD, CHD and stroke respectively<sup>25</sup>. The dose-response analysis suggested that the 'optimum' cheese consumption is about 40g/day.

A lower risk of CHD was observed in a newly published meta-analysis comparing the highest with the lowest category of cheese intake<sup>20</sup>. Analyses of dairy product consumption in 13 studies also found lower CVD for women (13% reduction in risk with high dairy intake) but not for men<sup>26</sup>. A systematic review of the association between dairy product consumption and risk of various cardiovascular-related clinical outcomes reports favourable

associations between intakes of total dairy, low-fat dairy, cheese, and fermented dairy and the risk of stroke<sup>19</sup>. Similarly, a recent dose-response meta-analysis combining data from 29 prospective cohort studies demonstrated neutral associations between dairy products and cardiovascular mortality<sup>18</sup>.

## Potential dairy matrix mechanisms

**The explanation for the finding that dairy foods, even those containing fat and saturated fat such as cheese, have a neutral or even a beneficial effect on CVD is likely to lie in the complex structural and nutritional composition of the dairy matrix<sup>27</sup>.**

Although some dairy foods contain saturated fatty acids, they are also rich in nutrients and bioactive components such as **calcium, potassium, phosphorus and bioactive peptides** that may modify CVD risk through, for example, positive effects on blood pressure, weight and diabetes. In addition, dairy constituents such as bioactive peptides may have direct effects on cardiovascular parameters including blood clotting, arterial stiffness endothelial function and inflammation<sup>28</sup>. This may help counter any negative effects of saturated fat in dairy on blood lipids and CVD risk.

**Moreover, interactions of the components and structure of the dairy matrix including calcium, phosphorus, the milk fat globule membrane and starter cultures, have been shown to modify saturated fatty acid-induced increases in blood lipids<sup>27, 29</sup>.** There is evidence that **calcium** in dairy foods, through its effects on binding fat and decreasing its absorption in the gut, may reduce the potential rise in LDL cholesterol following saturated fat consumption<sup>30-32</sup>. For example, cheese does not increase LDL cholesterol compared with butter of equal fat content<sup>31</sup>. Similarly, compared with a low-calcium control diet, milk- and cheese-based diets lessened saturated fatty acid-induced increases in total and LDL cholesterol<sup>33</sup>. It may be important for this beneficial effect that fat and calcium are embedded in the same food matrix, as is the case for milk and cheese<sup>27</sup>. In fact, a recent randomized controlled trial found that dairy fat, eaten in the form of cheese, appeared to affect blood lipids differently

compared with the same constituents eaten in separate matrices, with significantly lower total cholesterol observed when all nutrients were consumed within the cheese matrix<sup>29</sup>.

**Phosphorus** in the dairy matrix may also interact with calcium to influence blood lipids; calcium phosphate binds bile acids and fatty acids, and increases their excretion<sup>34</sup>. It has been suggested too that the membrane which encloses milk fat (the **milk fat globule membrane**; MFGM) and which is rich in bioactive phospholipids and proteins may have a beneficial role in modulating blood lipids<sup>35, 36</sup>. Fermented dairy foods may also modify blood lipids through favouring gut bacteria which produce short-chain fatty acids (SCFA), and which in turn have a positive effect on lipids<sup>37</sup>. The structure and texture of dairy food matrices can also influence digestion kinetics and uptake of lipids; the physical structure of cheese, for example, can slow fat digestion<sup>38</sup>.

**It is also increasingly recognised that individual saturated fatty acids have different effects on blood lipids;** several of those in milk fat do not have an adverse effect on LDL ('bad') cholesterol or other markers of CVD risk including HDL ('good') cholesterol and the ratio of total to HDL cholesterol<sup>39</sup>. In addition, the total fatty acid profile of a food, not just its saturated fatty acid content, is important. Milk fat includes a number of fatty acids which may have beneficial effects on CVD risk factors such as blood lipids and markers of inflammation. These include conjugated linoleic acid (cis-9, trans-11 CLA) and trans palmitoleic acid (trans-C16:1)<sup>40-42</sup>.

**It is evident that in terms of the effects of milk and dairy foods on cardiovascular health, the whole food, and the dietary pattern, rather than an individual component such as saturated fat should be taken into account. In this respect, the weight of evidence suggests no adverse effects of regularly consuming milk and dairy foods on cardiovascular health and rather, in some studies, a cardio-protective effect has been observed.**

## References

1. Elwood PC et al. Milk drinking, ischaemic heart disease and ischaemic stroke II. Evidence from cohort studies. *Eur J Clin Nutr.* 2004; 58: 718-724.
2. Goldbohm RA et al. Dairy consumption 10-y total and cardiovascular mortality: a prospective cohort study in the Netherlands. *Am J Clin Nutr.* 2011; 93: 615-627.
3. van Aerde MA et al. Dairy intake in relation to cardiovascular disease mortality and all-cause mortality: the Hoorn Study. *Eur J Nutr.* 2013; 52: 609-616.
4. Praagman J et al. Dairy products and the risk of stroke and coronary heart disease the Rotterdam Study. *Eur J Nutr.* 2015; 54: 981-990.
5. Dalmeijer GW et al. Dairy intake and coronary heart disease or stroke—a population-based cohort study. *Int J Cardiol.* 2013; 167: 925-929.
6. Larsson SC et al. Dairy consumption and risk of stroke in Swedish women and men. *Stroke.* 2012; 43: 1775-1780.
7. Sonestedt E et al. Dairy products and its association with incidence of cardiovascular disease: the Malmö diet and cancer cohort. *Eur J Epidemiol.* 2011; 26: 609-618.
8. Michaëlsson K et al. Milk intake and risk of mortality and fractures in women and men: cohort studies. *BMJ.* 2014; 349: g6015.
9. Hellstrand S. The statistical analysis and reality. Letter to *BMJ.* *BMJ.* 2014; 349: g6015.
10. Bergholdt HK et al. Milk intake is not associated with low risk of diabetes or overweight-obesity: a Mendelian randomization study in 97,811 Danish individuals. *Am J Clin Nutr.* 2015; 102: 487-496.
11. Bongard V et al. Association of dietary patterns with 14-year all-cause mortality and cause-specific mortality. *Eur Heart J.* 2012; 33 (S1): 609-610.
12. Sellem, L et al. Consumption of dairy products and cardiovascular disease risk: Results from the French prospective cohort NutriNet-Santé. *Proc Nut Soc.* 2020; 79(OCE2), E152. doi:10.1017/S0029665120001007
13. Dehghan, M et al. Association of dairy intake with cardiovascular disease and mortality in 21 countries from five continents (PURE): A prospective cohort study. *Lancet* 201; 392: 2288–2297.
14. Menté A et al. A systematic review of the evidence supporting a causal link between dietary factors and coronary heart disease. *Arch Intern Med.* 2009; 169: 659-669.
15. Elwood PC et al. The consumption of milk and dairy foods and the incidence of vascular disease and diabetes: an overview of the evidence. *Lipids.* 2010; 45: 925-939.
16. Soedamah-Muthu SS et al. Milk and dairy consumption and incidence of cardiovascular diseases and all-cause mortality: dose-response meta-analysis of prospective cohort studies. *Am J Clin Nutr.* 2011; 93: 158-171.
17. Larson SC et al. Milk consumption and mortality from all causes, cardiovascular disease, and cancer: a systematic review and meta-analysis. *Nutrients.* 2015; 7: 7749-7763.
18. Guo J et al. Milk and dairy consumption and risk of cardiovascular diseases and all-cause mortality: dose-response meta-analysis of prospective cohort studies. *Eur J Epidemiol* 2017; 32: 269–287.
19. Drouin-Chartier J-P et al. Systematic review of the association between dairy product consumption and risk of cardiovascular-related clinical outcomes. *Adv Nutr.* 2016; 7: 1026–1040.
20. Jakobsen MU et al. Intake of dairy products and associations with major atherosclerotic cardiovascular diseases: a systematic review and meta-analysis of cohort studies. *Sci Rep.* 2021; 11: 1303.
21. Qin LQ et al. Dairy consumption and risk of cardiovascular disease: an updated meta-analysis of prospective cohort studies. *Asia Pac J Clin Nutr.* 2015; 24: 90-100.
22. Hu D et al. Dairy foods and risk of stroke: a meta-analysis of prospective cohort studies. *Nutr Metab Cardiovasc Dis.* 2014; 24: 460-469.
23. de Goede J et al. Dairy consumption and risk of stroke: a systematic review and updated dose-response meta-analysis of prospective cohort studies. *J Am Heart Assoc.* 2016 May 20; 5(5). doi: 10.1161/JAHA.115.002787.
24. Alexander D et al. Dairy consumption and CVD: a systematic review and meta-analysis. *Br J Nutr.* 2016; 115: 737–750.
25. Chen GC et al. Cheese consumption and risk of cardiovascular disease: a meta-analysis of prospective studies. *Eur J Nutr* 2017; 56: 2565-2575.
26. Mishali M et al. Association between dairy intake and the risk of contracting type 2 diabetes and cardiovascular diseases: a systematic review and meta-analysis with subgroup analysis of men versus women. *Nutr Rev.* 2019; 77: 417–429.

27. Thorning TK et al. Whole dairy matrix or single nutrients in assessment of health effects: current evidence and knowledge gaps. *Am J Clin Nutr* 2017; 105:1–13.
28. Ricci-Cabello et al. Possible role of milk-derived bioactive peptides in the treatment and prevention of metabolic syndrome. *Nutr Rev*. 2012; 70: 241-255.
29. Feeney, EL et al. Dairy matrix effects: Response to consumption of dairy fat differs when eaten within the cheese matrix - A randomized controlled trial. *Am J Clin Nutr*. 2018; 108: 667–674.
30. Lorenzen JK & Astrup A. Dairy calcium intake modifies responsiveness of fat metabolism and blood lipids to a high-fat diet. *Br J Nutr*. 2011; 105: 1823-1831.
31. Soerensen KV et al. Effect of dairy calcium from cheese and milk on fecal fat excretion, blood lipids, and appetite in young men. *Am J Clin Nutr*. 2014; 99: 984-991.
32. Lorenzen JK et al. Milk minerals modify the effect of fat intake on serum lipid profile: results from an animal and a human short-term study. *Br J Nutr*. 2014; 111: 1412-1420.
33. Hjerpsted J et al. Cheese intake in large amounts lowers LDL-cholesterol concentrations compared with butter intake of equal fat content. *Am J Clin Nutr*. 2011; 94: 1479-1484.
34. Govers MJ et al. Characterization of the adsorption of conjugated and unconjugated bile acids to insoluble, amorphous calcium phosphate. *J Lipid Res*. 1994; 35: 741–748.
35. Rosqvist F et al. Potential role of milk fat globule membrane in modulating plasma lipoproteins, gene expression, and cholesterol metabolism in humans: a randomized study. *Am J Clin Nutr*. 2015; 102: 20-30.
36. Anto L et al. Milk polar lipids: underappreciated lipids with emerging health benefits. *Nutrients*. 2020;12(4):1001. doi: 10.3390/nu12041001.
37. St-Onge MP et al. Consumption of fermented and nonfermented dairy products: effects on cholesterol concentrations and metabolism. *Am J Clin Nutr*. 2000; 71: 674–681.
38. Schmidt JM et al. Influence of type of dairy matrix micro- and macrostructure on in vitro lipid digestion. *Food Funct* 2020; 11: 4960 – 4972.
39. Kris-Etherton PM & Yu S. Individual fatty acid effects on plasma lipids and lipoproteins: human studies. *Am J Clin Nutr*. 1997; 65: 1628S-1644S.
40. Tricon S et al. Opposing effects of cis-9, trans-11 and trans-10, cis-12 conjugated linoleic acid on blood lipids in healthy humans. *Am J Clin Nutr*. 2004; 80: 614-620.
41. Mozaffarian D et al. Trans-palmitoleic acid, metabolic risk factors, and new-onset diabetes in US adults: a cohort study. *Ann Intern Med*. 2010; 153: 790-799.
42. Kratz et al. The relationship between high-fat dairy consumption and obesity, cardiovascular, and metabolic disease. *Eur J Nutr*. 2013; 52: 1-24.45.



by



**European**  
**Milk**  
**Forum**

[www.milknutritiousbynature.eu](http://www.milknutritiousbynature.eu)